

Package ‘symmetry’

August 24, 2019

Title Testing for Symmetry of Data and Model Residuals

Version 0.1.0

Author Blagoje Ivanović [aut, cre]

Bojana Milošević[aut]

Marko Obradović [aut]

Date 2019-08-24

Maintainer Blagoje Ivanović <blagoje_ivanovic@matf.bg.ac.rs>

Description Implementations of a large number of tests for symmetry and their bootstrap variants, which can be used for testing the symmetry of IID samples around a known or unknown mean. Functions are also there for testing the symmetry of model residuals around zero. Currently, the supported models are linear models and GARCH models (fitted with the 'fGarch' package). All tests are implemented using the 'Rcpp' package which ensures great performance of the code.

Depends R (>= 3.1.0)

License MIT + file LICENSE

Encoding UTF-8

LazyData true

Imports Rcpp, Rdpack

RdMacros Rdpack

LinkingTo Rcpp, RcppArmadillo

RoxygenNote 6.1.1

SystemRequirements C++11

Suggests knitr, rmarkdown, sn, fGarch, testthat

Roxygen list(markdown = TRUE)

R topics documented:

rmixnorm	2
rsl	2

symmetry	3
symmetry_test	3
TestStatistics	6

Index	9
--------------	----------

rmixnorm	<i>Mixture of 2 normal distributions</i>
----------	--

Description

Generates random numbers from a mixture of 2 normal distributions

Usage

```
rmixnorm(n, mean1 = 0, sd1 = 1, mean2 = 0, sd2 = 1, p = 0.5)
```

Arguments

n	number of observations
mean1	mean of the first normal
sd1	standard deviation of the first normal
mean2	mean of the second normal
sd2	standard deviation of the second normal
p	probability of the first normal

Value

Vector of random numbers from the specified mixture of normals.

rs1	<i>Azzalini skew logistic distribution</i>
-----	--

Description

Generates random numbers from the skew logistic distribution

Usage

```
rs1(n = 1, xi = 0, omega = 1, alpha = 0, dp = NULL)
```

Arguments

n	sample size.
xi	vector of location parameters.
omega	vector of (positive) scale parameters.
alpha	vector of slant parameters.
dp	a vector of length 3 whose elements represent the parameters described above. If dp is specified, the individual parameters cannot be set.

Value

Vector of random numbers from Azzalini skew logistic distribution.

symmetry	<i>symmetry: A package which implements tests for symmetry of IID data, linear models and GARCH models</i>
----------	--

Description

The package contains a large number of tests for symmetry (and their bootstrap variants), which can be used to test the symmetry of IID samples or of model residuals. Currently, the supported models are linear models and GARCH models (fitted with the 'fGarch' package). The tests are implemented using Rcpp which ensures great performance.

Details

To see the available tests, see [TestStatistics](#)

For documentation on how to perform the tests, see [symmetry_test](#)

symmetry_test	<i>Perform symmetry tests</i>
---------------	-------------------------------

Description

This is a generic function used to perform symmetry tests on numeric vectors or objects of class lm (linear models) and objects of class fGARCH (GARCH models fitted with the fGarch package).

Usage

```

symmetry_test(x, ...)

## Default S3 method:
symmetry_test(x, stat, mu = NULL, bootstrap = TRUE,
  B = 1000, boot_method = c("sign", "reflect"), trim = 0, k = 0,
  ...)

## S3 method for class 'lm'
symmetry_test(x, stat, B = 1000, boot_method = c("sign",
  "reflect"), k = 0, ...)

## S3 method for class 'fGARCH'
symmetry_test(x, stat, B = 1000, burn = 0,
  boot_method = c("sign", "reflect"), k = 0, approximate = FALSE,
  ...)

```

Arguments

x	an object of class numeric, lm or fGARCH
...	not used
stat	a character vector indicating the test statistic to be used (see Available Test Statistics)
mu	the centre parameter around which to test symmetry
bootstrap	a logical indicating whether to use bootstrap
B	the number of bootstrap replications
boot_method	the method of bootstrap sample generation (see Details)
trim	the trim value used for estimating the centre (as used in "mean")
k	the k parameter of the statistic, ignored if the test statistic doesn't depend on a parameter (see Test Statistics)
burn	the number of elements to remove from the beginning of the time series for testing
approximate	a logical indicating whether to use the faster approximate bootstrap method (see Details)

Details

The tests are performed using bootstrap procedures or using asymptotic results, where applicable. Currently, two methods of generating a bootstrap sample from the null distribution are available. The "sign" method generates the bootstrap sample by multiplying the existing sample by -1 or 1 at random (with equal probabilities), essentially randomizing the sign of the data, giving a symmetric distribution. The "reflect" method reflects the sample around zero and samples `length(x)` elements with replacement. In practice, it has been shown that the "sign" method is almost always better, thus is the default.

For numeric data, the tests can be performed around a known (parameter "mu") or unknown centre. When the centre is known, the bootstrap method gives the same results as a Monte Carlo simulation of the p value, for tests which are distribution free. For unknown centre (when mu = NULL), bootstrap must be used and the estimate of the centre used is the trimmed mean, with trim parameter "trim". By default, the mean is taken (trim = 0).

For linear models, the tests are based on a bootstrap procedure as in (Allison and Pretorius 2017) and are used to test the symmetry of the residuals around zero.

For GARCH models (must be fitted with the 'fGarch' package), the tests are also based on bootstrap and test for symmetry of the residuals around zero. An approximation of the bootstrap procedure is available where the residuals are treated as iid data, which is much faster and has been shown to give similar results to the default bootstrap procedure (described in (Klar et al. 2012)).

For a comparison of the performance of various tests of symmetry around an unknown centre, see (Milošević and Obradović 2019)).

Value

An object of class "htest" containing the results of the testing.

References

Allison J, Pretorius C (2017). "A Monte Carlo evaluation of the performance of two new tests for symmetry." *Computational Statistics*, **32**(4), 1323–1338. doi: [10.1007/s0018001606804](https://doi.org/10.1007/s0018001606804).

Klar B, Lindner FL, Meintanis S (2012). "Specification tests for the error distribution in GARCH models." *Computational Statistics & Data Analysis*, **56**(11), 3587–3598. doi: [10.1016/j.csda.2010.05.029](https://doi.org/10.1016/j.csda.2010.05.029).

Milošević B, Obradović M (2019). "Comparison of efficiencies of some symmetry tests around an unknown centre." *Statistics*, **53**(1), 43–57.

Examples

```
set.seed(1)

# IID samples
x <- rnorm(50)
symmetry_test(x, "MOI", bootstrap = FALSE, k = 3, mu = 0)
symmetry_test(x, "MOI", bootstrap = TRUE, k = 3, mu = 0)
symmetry_test(x, "MOI", bootstrap = TRUE, k = 3)
x <- rsl(50, alpha = 1.5)
symmetry_test(x, "MOI", bootstrap = FALSE, k = 3, mu = 0)
symmetry_test(x, "MOI", bootstrap = TRUE, k = 3, mu = 0)
symmetry_test(x, "MOI", bootstrap = TRUE, k = 3)

# Linear models
lin_model <- lm(dist ~ speed, cars)
symmetry_test(lin_model, "B1")

# Garch models
library(fGarch)
specskew19 = fGarch::garchSpec(model = list(omega = 0.1,
```

```

                                alpha = 0.3,
                                beta = 0.3,
                                skew = 1.9),
                                cond.dist = "snorm")

x <- fGarch::garchSim(specskew19, n = 500)
g <- fGarch::garchFit(~garch(1,1), x, cond.dist = "QMLE",
                      include.mean = FALSE, trace = FALSE)
symmetry_test(g, "CH", B=400, burn = 100) # slower
symmetry_test(g, "CH", B=400, burn = 100, approximate = TRUE)

```

TestStatistics

Available test statistics for symmetry tests

Description

The list of implemented test statistics and their functions

Usage

B1(X)

BHI(X)

BHK(X)

CH(X)

CM(X)

K2U(X)

K2(X)

KS(X)

SGN(X)

WCX(X)

MGG(X)

MOI(X, k)

MOK(X, k)

M(X)

NAI(X, k)

NAK(X, k)

S1(X, k)

S2(X, k)

T1(X, k)

T2(X, k)

Arguments

X	the numeric vector for which to calculate the test statistic
k	the 'k' parameter in the formula (if applicable)

Details

Below is a list of the implemented test statistics in the package. Each statistic is listed by its name, a code string (e.g. 'B1', 'CM', 'MOI') and the formula of the statistic which is evaluated. The code string is used as an argument to the [symmetry_test](#) function. Some statistics depend on a parameter 'k' which can be seen from the formulas and is also passed as an argument.

Each statistic is implemented as a function with the same name as the code string, so the name of the function is passed as the argument "stat" to the [symmetry_test](#) function

Value

The value of the test statistic.

Test statistics

The list of available statistics in the format "code(s) : name (reference)"

- M : The Mira test statistic (see (Mira 1999))
- CM : The Cabilio–Masaro test statistic (see (Cabilio and Masaro 1996))
- MGG : The Miao, Gel and Gastwirth test statistic (see (Miao et al. 2006))
- B1 : The $\sqrt{b_1}$ test statistic (see (Milošević and Obradović 2019))
- KS : The Kolmogorov–Smirnov test statistic (see (Milošević and Obradović 2019))
- SGN : The Sign test statistic (see (Milošević and Obradović 2019))
- KS : The Wilcoxon test statistic (see (Milošević and Obradović 2019))
- CH : The characterization based test statistic (see (Feuerverger et al. 1977))
- BHI : The Litvinova test statistic (see (Litvinova 2001))
- BHK : The Baringhaus and Henze test statistic (see (Baringhaus and Henze 1992))

- MOI and MOK : The Milošević and Obradović test statistics (see (Milošević and Obradović 2016))
- NAI and NAK : The Nikitin and Ahsanullah test statistics (see (Nikitin and Ahsanullah 2015))
- K2 and K2U : The Božin, Milošević, Nikitin and Obradović Kolmogorov type statistics based on V- and U- statistics respectively (see (Božin et al. 2018))
- T1, T2, S1 and S2 : The Allison and Pretorius test statistics (see (Allison and Pretorius 2017))

References

- Allison J, Pretorius C (2017). “A Monte Carlo evaluation of the performance of two new tests for symmetry.” *Computational Statistics*, **32**(4), 1323–1338. doi: [10.1007/s0018001606804](https://doi.org/10.1007/s0018001606804).
- Baringhaus L, Henze N (1992). “A characterization of and new consistent tests for symmetry.” *Communications in statistics-theory and methods*, **21**(6), 1555–1566. doi: [10.1080/03610929208830863](https://doi.org/10.1080/03610929208830863).
- Božin V, Milošević B, Nikitin Y, Obradović M (2018). “New Characterization-Based Symmetry Tests.” *Bulletin of the Malaysian Mathematical Sciences Society*, 10–1007. doi: [10.1007/s40840-01806803](https://doi.org/10.1007/s40840-01806803).
- Cabilio P, Masaro J (1996). “A simple test of symmetry about an unknown median.” *Canadian Journal of Statistics*, **24**(3), 349–361. doi: [10.2307/3315744](https://doi.org/10.2307/3315744).
- Feuerverger A, Mureika R, others (1977). “The empirical characteristic function and its applications.” *The Annals of Statistics*, **5**(1), 88–97. doi: [10.1214/aos/1176343742](https://doi.org/10.1214/aos/1176343742).
- Litvinova V (2001). “New nonparametric test for symmetry and its asymptotic efficiency.” *Vestnik St. Petersburg University Mathematics*, **34**(4), 12–14.
- Miao W, Gel YR, Gastwirth JL (2006). “A new test of symmetry about an unknown median.” In *Random Walk, Sequential Analysis And Related Topics: A Festschrift in Honor of Yuan-Shih Chow*, 199–214. World Scientific.
- Milošević B, Obradović M (2019). “Comparison of efficiencies of some symmetry tests around an unknown centre.” *Statistics*, **53**(1), 43–57.
- Milošević B, Obradović M (2016). “Characterization based symmetry tests and their asymptotic efficiencies.” *Statistics & Probability Letters*, **119**, 155–162.
- Mira A (1999). “Distribution-free test for symmetry based on Bonferroni’s measure.” *Journal of Applied Statistics*, **26**(8), 959–972. doi: [10.1080/02664769921963](https://doi.org/10.1080/02664769921963).
- Nikitin Y, Ahsanullah M (2015). “New U-empirical Tests of Symmetry Based on Extremal Order Statistics, and their Efficiencies.” In *Mathematical Statistics and Limit Theorems*, 231–248. Springer. doi: [10.1007/9783319124421_13](https://doi.org/10.1007/9783319124421_13).

Index

Available Test Statistics, [4](#)

B1 (TestStatistics), [6](#)
BHI (TestStatistics), [6](#)
BHK (TestStatistics), [6](#)

CH (TestStatistics), [6](#)
CM (TestStatistics), [6](#)

K2 (TestStatistics), [6](#)
K2U (TestStatistics), [6](#)
KS (TestStatistics), [6](#)

M (TestStatistics), [6](#)
MGG (TestStatistics), [6](#)
MOI (TestStatistics), [6](#)
MOK (TestStatistics), [6](#)

NAI (TestStatistics), [6](#)
NAK (TestStatistics), [6](#)

rmixnorm, [2](#)
rsl, [2](#)

S1 (TestStatistics), [6](#)
S2 (TestStatistics), [6](#)
SGN (TestStatistics), [6](#)
symmetry, [3](#)
symmetry-package (symmetry), [3](#)
symmetry_test, [3](#), [3](#), [7](#)

T1 (TestStatistics), [6](#)
T2 (TestStatistics), [6](#)
Test Statistics, [4](#)
TestStatistics, [3](#), [6](#)

WCX (TestStatistics), [6](#)